



上海中醫藥大學

Shanghai University of Traditional Chinese Medicine


2014

GB/T7713-1987

GB/T 7714-2015

GB/T7713.1-2006

1.1	1
1.1.1	1
1.1.2	1
1.1.3	1
1.1.4	1
1.1.5	1
1.1.6	1
1.2	2
1.3	2
1.4	2
1.5	3
1.6	3
1.6.1	4
1.6.2	4
1.7	4
1.7.1	5
1.7.2	5
1.7.3	5
1.7.4	6
1.8	8
1.8.1	8
1.8.2	11
1.8.3	12
1.9	12
1.10	12
21	14
22	14



--	--	--	--

•

•

•

•

•

•

表 3 非浸润性膀胱癌不同中医证型患者生存质量评分比较($\bar{x}\pm s$, 分)

证型	n	乏力	疼痛	呼吸困难	失眠	食欲丧失	便秘	排尿症状	肠道症状
瘀毒蕴结型	144	64.4±22.7	69.7±23.4*	5.9±15.3	44.3±24.5	40.9±27.6	55.5±18.3*	52.9±19.6*	39.5±18.8*
湿热下注型	149	64.3±23.5	57.6±21.3*	5.8±15.8	45.1±23.4	42.7±25.1	23.4±8.1	68.8±16.7*	58.4±18.3*
脾肾两虚型	18	67.5±22.8	52.8±20.7*	6.4±15.2	45.6±22.9	57.5±23.4*	47.9±22.5*	46.2±16.5*	47.1±17.9*
无证可辨/其他证型	31	43.0±22.8*	40.4±23.4*	6.1±15.8	43.3±24.5	41.1±23.5	39.1±24.7*	41.5±18.6*	36.1±15.0*

生活质量	证型	n	经济困难	躯体功能	认知功能	情绪功能	社会功能	总体健康状况	对将来担心	性
5±16.6	瘀毒蕴结型	144	57.6±31.5	60.1±18.4	78.3±20.1	41.7±24.6	35.4±22.7	21.6±14.3	51.6±10.3	7
8±13.4	湿热下注型	149	53.4±27.3	57.5±18.0	78.6±21.7	39.8±17.4	39.1±19.9	23.4±8.1	50.7±12.1	6
6±16.2	脾肾两虚型	18	52.1±28.8	58.2±18.1	79.4±21.3	38.4±17.2	32.5±21.0*	19.5±8.5*	51.5±11.8	6
9±11.5*	肝肾阴虚型	44	58.4±32.3	56.0±19.5	79.1±21.2	42.2±24.5	36.6±23.8	24.9±13.4	50.4±11.6	4
8±15.8	无证可辨/其他证型	31	56.4±31.3	65.3±18.2*	77.5±24.8	47.5±15.3*	40.8±24.3	40.3±6.5*	52.9±15.3	6

*与湿热下注型比较, $P < 0.05$; 与瘀毒蕴结型比较, $P < 0.05$

Table 2 Treatment and the prognosis of recurrence breast cancer patients

Index	TNBC (%)	Non-TNBC (%)	χ^2	P
Surgery type				
Total mastectomy	392(86.15)	230(85.74)		
No surgery / unknown	11(2.42)	94(3.50)	1.531	0.4
Neoadjuvant chemotherapy				
No	330(72.53)	2067(76.95)		
Yes	110(24.18)	553(20.59)	4.435	0.1
Unknown	15(3.30)	66(2.46)		
Adjuvant radiotherapy				
No	232(50.99)	1462(54.43)		
Yes	208(45.71)	1116(41.55)	2.977	0.2
Unknown	15(3.30)	108(4.02)		
Adjuvant chemotherapy				
No	25(5.49)	230(8.56)		
Yes	422(92.75)	2359(87.83)	9.574	0.0
Unknown	8(1.76)	97(3.61)		
DFS(months)				
≤24	268(58.90)	1096(40.80)	51.867	<0.001
>24	187(41.10)	1590(59.20)		
Type of recurrence				
Distant metastasis	357(78.46)	2112(78.63)		
Local recurrence	89(19.56)	486(18.09)	2.582	0.275
Unknown	9(1.98)	88(3.28)		
Liver metastasis				
Yes	71(15.60)	562(20.92)		
No	384(84.40)	2117(78.82)	8.169	0.017
Unknown	0	7(0.26)		
Lung metastasis				
Yes	161(35.29)	704(25.82)		
No	294(64.62)	1885(70.18)	7.262	0.026
Unknown	0	7(0.26)		
Bone metastasis				
Yes	129(28.35)	1041(38.76)		
No	326(71.65)	1638(60.98)	19.573	<0.001
Unknown	0	7(0.26)		
Brain metastases				
Yes	17(3.74)	84(3.13)		
No	438(96.26)	2595(96.61)	1.639	0.441
Unknown	0	7(0.26)		

表 1 两组患者临床资料比较 [例 (%) $\bar{x}\pm s$]

临床资料	异丙托溴铵组 (96例)	安慰剂组 (94例)	整体人群 (190例)
性别			
男	83 (86.5)	85 (90.4)	168 (88.4)
女	13 (13.5)	9 (9.6)	22 (11.6)
年龄(岁)	63.4±6.5	62.4±6.4	62.9±6.5
年龄≥60岁	67 (69.8)	65 (69.1)	132 (69.5)
急性支气管扩张剂反应性			
是	13 (13.5)	13 (13.8)	26 (13.7)
否	83 (86.5)	81 (86.2)	164 (86.3)
身高 (cm)	165.90±7.30	168.60±7.90	167.20±7.70
体重 (kg)	61.80±11.58	64.07±11.66	62.93±11.65
体重指数 (kg/m ²)	22.38±3.42	22.50±3.51	22.44±3.46
吸烟状况			
从不吸烟	15 (15.6)	6 (6.4)	21 (11.1)
以前吸烟	55 (57.3)	54 (57.4)	109 (57.4)
目前吸烟	26 (27.1)	34 (36.2)	60 (31.6)
饮酒史			
不饮酒者	60 (62.5)	54 (57.4)	114 (60.0)
饮酒者	36 (37.5)	40 (42.6)	76 (40.0)



示例如下：



示例如下：



示例如下：

示例如下：




示例如下：



示例如下：







-
-
-
-
-
-
-
-
-
-
-